

**Draft**  
**Bull Trout and Dolly Varden  
Management Plan**

Washington Department of Fish and Wildlife  
Fish Management Program  
600 Capitol Way North  
Olympia, WA 98501-1091

# Table of Contents

---

Executive Summary .....	ii
Introduction .....	1
Management Plan Goal .....	1
Taxonomy .....	2
Life History and Habitat Requirements .....	2
Distribution and Stock Status .....	3
Population Maintenance .....	4
Fisheries Management .....	6
Habitat Maintenance .....	6
Enforcement .....	7
Research Monitoring and Evaluation .....	8
Literature Cited .....	11
Glossary .....	14

This work was funded in part by the Federal Aid in Sport Fish Restoration Act administered by the U.S. Fish and Wildlife Service.
---

# Executive Summary

---

The Bull Trout and Dolly Varden Management Plan describes the goal, objectives and strategies that will be used by the Washington Department of Fish and Wildlife (Department) for managing bull trout and Dolly Varden in Washington. The plan does not represent a consensus of all entities managing native char. However, the Department will work with Indian tribes and other management agencies to reach consensus through negotiations when implementing this plan at the local watershed level.

The goal of the Department is to restore and maintain the health and diversity of bull trout (*Salvelinus confluentus*) and Dolly Varden (*S. malma*) stocks and their habitats and allow recreational utilization within resource protection guidelines. This goal addresses the Department's mandate, the Fish and Wildlife Commission's policies, and the Wild Salmonid Policy. The intent of the goal is to address stock health beyond numerical abundance by ensuring the long-term productive capacity of bull trout and Dolly Varden and their habitats are maintained.

In the state of Washington, bull trout and Dolly Varden, two closely-related char species co-exist. They have considerable biological similarities with respect to life history and habitat requirements. For these reasons the goal, objectives and strategies in this plan address both bull trout and Dolly Varden in Washington.

The Department completed a stock status report for bull trout and Dolly Varden in 1997 which identified 80 stocks statewide. Sixteen percent were identified as healthy, 3% depressed, 6% critical and 75% as unknown.

This plan defines population maintenance, fisheries management and habitat maintenance objectives. In addition it describes the enforcement, monitoring, evaluation and research efforts needed to meet the bull trout and Dolly Varden management goal and objectives.

# Introduction

---

The Washington Fish and Wildlife Commission (Commission) is directed by the Washington State Legislature (RCW 77.04.055) to establish policies to preserve, protect and perpetuate fish and fish habitat and to maximize recreational fishing opportunities compatible with healthy and diverse fish populations. In 1995 the Commission established goals, policies and objectives for the Commission and the Washington Department of Fish and Wildlife (Department) to meet this mandate (Washington Fish and Wildlife Commission 1995).

In 1984 the Game Commission adopted a basic fishery management strategy (BSMS) for resident and anadromous trout in streams of the state of Washington (Washington Department of Game 1984) which in part focused management of bull trout and Dolly Varden (native char) in streams on sustainable wild productions.

The Department completed a draft bull trout/Dolly Varden Management Plan in 1992 (Washington Department of Wildlife 1992) and initiated a programmatic Environmental Impact Statement (EIS) for the management plan under the State Environmental Policy Act. Based in part on comments received during review of the preliminary draft EIS, the Department elected to postpone completion of the bull trout/Dolly Varden management plan to ensure it was consistent with the policy for wild salmonids being developed.

In 1997 the Commission adopted the Wild Salmonid Policy (WSP) containing goals and policies concerning wild salmonids agreed to provisionally by Western Washington Treaty Tribes (Washington Fish and Wildlife Commission 1997a), and additional policy guidance for Department staff (Washington Fish and Wildlife Commission 1997b). These documents address issues including harvest by fisheries, hatcheries management, and habitat protection needed to manage salmonid resources in Washington.

This management plan for bull trout and Dolly Varden in Washington State is consistent with the Commission goals, policies and objectives and provides specific guidance for management of these species by the Department under the WSP. This plan does not represent a consensus of all entities managing native char. However, the Department will work with Indian tribes and other management agencies to reach consensus through negotiations when implementing this plan at the local watershed level.

## Management Plan Goal

<p><b>To restore and maintain the health and diversity of bull trout and Dolly Varden stocks and their habitats and allow recreational utilization within resource protection guidelines.</b></p>
---

The intent of the goal is to address stock health beyond numerical abundance by ensuring the long-term productive capacity of bull trout and Dolly Varden and their habitats are maintained. The highest priority for management of native char will be resource protection. The specific objectives and strategies in this plan are grouped into several elements including population maintenance, fisheries management, and habitat

maintenance. In addition it describes the enforcement, monitoring, evaluation and research efforts needed to meet the bull trout and Dolly Varden management goal and objectives.

## Taxonomy

Species of the genus *Salvelinus* are trout-like in appearance, but generally have finer scales and exhibit a reverse coloration with light spotting on a darker background, while the true trout and salmon have black spots on a lighter background. Bull trout and Dolly Varden are western North America's most southern derivatives of an evolutionary parent *Salvelinus*, possibly the Arctic char (Morton, 1970; Behnke, 1980; Savvaitova, 1980; Balon, 1984).

In the past, coastal native char were considered Dolly Varden and native char in inland waters were considered bull trout. Cavender (1978) demonstrated that bull trout and Dolly Varden were separate species and believed the geographic range of bull trout and Dolly Varden overlapped in the Puget Sound area of Washington and along the British Columbia coast. Researchers using morphometric techniques have found no clear separation of the two species in Washington and that bull trout or a morphologically intermediate, or hybrid form predominated, even in coastal waters (Johnson and Mongillo, 1991; Kraemer, 1991). However, Crane et al. (1994) and others (reviewed by Utter 1994) demonstrated that genetic data can differentiate the two species. Use of genetic methods has recently documented the presence of pure Dolly Varden in Washington, that pure bull trout are present both east and west of the Cascade crest, and that the two species occur sympatrically in at least one river system in Western Washington (Leary and Allendorf 1997, Spruell and Allendorf 1997). Baxter et al. (1997) using molecular analyses confirmed the presence of natural hybrids and backcrosses between bull trout and Dolly Varden in the Thutade Lake drainage in north central British Columbia, and concluded that despite some apparent introgression, the two species appear to be maintaining distinct genomes.

## Life History and Habitat Requirements

Bull trout and Dolly Varden exhibit four life histories: anadromous, adfluvial, fluvial, and resident. These four survival strategies are common in various forms of Arctic char evolutionary derivatives and stocks worldwide. The four life history forms, or ecotypes, have unique life history and habitat requirement characteristics. However, there is a high degree of uncertainty regarding interrelationships, genetic and behavioral, among all four ecotypes.

**Anadromous:** Spawning and early rearing occurs in streams with major growth and maturation occurring in salt water. Stocks make frequent migrations into and out of lower mainstem rivers and estuaries. They often make long spawning migrations beginning early in summer. Mature adults normally range between 18 and 30 inches in total length.

**Adfluvial:** Spawning and rearing occurs in streams, but most growth and maturation occurs in lakes or reservoirs. Adults enter mainstem rivers early in summer, often holding near tributaries for months. Most mature adults range in size between 20 and 32 inches.

**Fluvial:** Spawning and early rearing occurs in smaller tributaries with major growth and maturation occurring in mainstem rivers. They may move randomly throughout river systems, generally congregating near spawning tributaries in the summer. Mature adults are usually smaller than anadromous or adfluvial char, ranging from 16 to 26 inches long.

**Resident:** All life stages (spawning, rearing, growth, maturation) occurs in small headwater streams, often upstream of impassable barriers. Mature adults can vary from 8 to 15 inches, but they are seldom larger than 12 inches in total length. Resident native char have been observed to mix and interbreed with migratory forms unless physically separated by barriers.

The native char of Washington evolved as an apex predator in most waters. Slow juvenile growth delays maturation until about age five, and reproduction may only occur on alternate years. Native char will live for 12 or more years in Washington, reaching sizes over 20 pounds where adequate forage is available. Stock densities of native char are generally much lower than other native game fish such as cutthroat trout (*Oncorhynchus clarki*) and rainbow trout (*O. mykiss*), or mountain whitefish (*Prosopium williamsoni*). The migratory forms of native char may travel long distances to reach spawning tributaries. Mature native char normally penetrate farther upstream than any other salmonids present in the watershed. Where native char and introduced brook trout (*Salvelinus fontinalis*) co-exist, hybridization may lead to eventual loss of native char stocks. A detailed discussion of the zoogeography and life history and habitat requirements of bull trout and Dolly Varden is presented by Brown (1992).

## Distribution And Stock Status

Bull trout or Dolly Varden are distributed widely in Washington State with reduced distribution from historical levels confined mainly to eastern Washington (Mongillo 1993). In addition to a reduction from historical numbers, several stocks in both eastern and western Washington have become fragmented where lower rivers are no longer utilized or free movement has been restricted. Relatively few native char are now observed in the mainstem Columbia and Snake Rivers. Native char have apparently been extirpated from the Chelan, lower Yakima, and Okanogan Basins. A number of stocks have been isolated above dams in river systems where they once roamed freely. However, with only a few exceptions, the overall geographic range of both continuous and disjunct stocks is believed similar to the historical range in Washington. Only one introduction of hatchery released native char has been verified in Washington. Anadromous Dolly Varden fry (14,500) from southeast Alaska were released into Lake Chelan in 1966. However, they were not observed in the lake after the initial introduction.

The Department recently completed a stock status inventory for both bull trout and Dolly Varden using the methodologies developed by the Department and Western Washington Treaty Tribe's for salmon and steelhead

(Washington Department of Fisheries et al., 1993). A separate inventory for each species was not performed due to the considerable biological similarities between the species and methodologies to reliably separate them where they coexist have only recently been developed and have not been widely applied. The stock status report identified 80 stocks statewide: sixteen percent were identified as healthy, 3% depressed, 6% critical and 75% as unknown (Table 1). For additional information on native char distribution and stock status refer to Washington Department of Fish and Wildlife (1997a).

**Table 1.** Regional and statewide summary of bull trout/Dolly Varden stock status.

	Healthy	Depressed	Critical	Unknown	Extinct
<b>Puget Sound</b>					
North Puget Sound	1	0	0	8	0
South Puget Sound	0	0	0	6	0
Hood Canal	1	0	0	2	0
Strait of Juan de Fuca	1	0	0	3	0
<b>Totals</b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>19</b>	<b>0</b>
<b>Coastal</b>					
North Coast	1	0	0	5	0
Grays Harbor	0	0	0	1	0
<b>Totals</b>	<b>1</b>	<b>0</b>	<b>0</b>	<b>6</b>	<b>0</b>
<b>Columbia River</b>					
Lower Columbia	0	1	0	0	0
Upper Columbia	9	1	5	35	0
<b>Totals</b>	<b>9</b>	<b>2</b>	<b>5</b>	<b>35</b>	<b>0</b>
Washington State					
80 Total Stocks	13	2	5	60	0
Percent of Total	16%	3%	6%	75%	0%

## Population Maintenance

The Department will use the stock (see glossary for definition) as the management unit for bull trout and Dolly Varden in Washington. Stock (population) maintenance will be achieved through strategies defined in Washington Department of Game (1984), summarized by Wright (1992) and described below. Bull trout and Dolly Varden stocks will be managed to ensure long-term surplus production, and to provide recreational harvest opportunities and ecological benefits consistent with the Fisheries Management and Habitat Maintenance objectives of this plan.

**Maintain and Restore Stock Distribution** - The Department will manage native char stocks and their habitat to promote distribution throughout their historic range.

*Implementation Strategy:*

- ! Habitat necessary for sustaining critical life history stages of native char including spawning and rearing will be protected or restored through efforts described in the habitat maintenance objectives.
- ! The Department will work through processes identified in the habitat maintenance objectives to protect current migratory corridors connecting remote headwater areas or restore historical corridors.

**Maintain and Restore Stock Abundance and Natural Biological Characteristics** - All stocks will be managed to maintain recruitment levels that ensure stable or increasing population densities and a natural age structure through maturation.

*Implementation Strategy:*

- ! Where habitat is limiting, implement measures to restore habitat necessary for sustaining critical life history stages of native char including spawning and rearing.
- ! Manage stocks with recreational fisheries to ensure through minimum size limits, that one full age class of mature females is allowed to spawn at least once prior to being subject to harvest. This will be defined as the youngest age class with a majority (more than 50%) of mature females.
- ! Sanctuaries and Refuges will be used to protect some stocks from most of the effects of degraded habitat, harvest and hatchery influences.

**Reestablish Stocks in Historically Inhabited Areas** - Stocks will be reestablished using supplementation in areas where they were found historically, where habitat is suitable to support healthy populations and no surviving pockets of native char are found.

*Implementation Strategy:*

- ! Stocks will be reestablished using supplementation beginning with pilot projects to test their feasibility in areas where they historically occurred, only in the context of a comprehensive recovery plan. Supplementation pilot projects will use guidance described in the Watershed Recovery Inventory Project Report (Washington Department of Fish and Wildlife 1997b).

**Conserve Genetic Diversity of Stocks** - Genetic diversity will be maintained within and among stocks to allow local adaptation to occur with changing environmental conditions over the long term.

*Implementation Strategy:*

- ! Habitat will be protected so that the distribution and amount of habitat is sufficient to maintain genetic diversity and promote local adaptation.



- ! Sanctuaries and Refuges will be used to protect some stocks from most of the effects of degraded habitat, harvest and hatchery influences.
- ! Where brook trout and native char currently overlap, the management emphasis will be to reduce or eliminate hybridization between them. This will generally be accomplished by using sport fishing regulations that allow higher harvest through increased daily limits, or with active suppression of brook trout.
- ! Brook trout will not be stocked into waters where there is a likelihood of interaction with native char stocks.

## **Fisheries Management**

Consistent with the population maintenance objectives, fishery management efforts generally emphasize resource protection of native char stocks by restricting recreational fishing and harvest.

Fisheries will be managed consistent with the use of the BSMS (Washington Department of Game 1984).

**Recreational Fisheries Will Only Be Allowed on Healthy Stocks with Surplus Production.**

*Implementation Strategy:*

- ! Directed sport fisheries will only be allowed on healthy stocks. Minimum size limits will be set based on life history type to ensure that a full age-class of females spawn at least once prior to recruitment into the fishery. Directed fisheries will not be allowed on stocks with depressed, critical or unknown status.
- ! Fishing closures for all species will be implemented in areas and times to protect critical spawning and rearing native char.
- ! Selective gear restrictions for all species will be implemented in areas and times to reduce incidental fishing mortality of native char.

## **Habitat Maintenance**

Many government agencies have jurisdiction for habitat protection and restoration in Washington State. The recently adopted WSP (Washington Fish and Wildlife Commission 1997a) identifies policies and performance measures for habitat maintenance that will be implemented by the Department for native char species. Habitat protection and restoration will occur primarily through a combination of locally-based watershed planning and general objectives that have the flexibility to implement performance measures and strategies based on local conditions.

This section summarizes habitat maintenance objectives and implementation strategies for native char to meet the performance measure and action strategies described in the WSP.

### **Maintain and Restore Habitat Necessary for Sustaining Critical Life History Stages Including Spawning, Rearing and Migration.**

#### *Implementation Strategy:*

- ! The Department, consistent with its authority of the State Hydraulic Code will incorporate conditions into Hydraulic Project Approvals (HPAs) to protect native char from negative impacts associated with activities that are subject to regulation under the HPA laws.
- ! The Department will provide input into State Environmental Policy Act review documents, forest practice applications, growth management plans, and water rights applications to protect native char habitat.
- ! The Department will actively participate on the Joint Cabinet for Natural Resources and will work with Tribes, local governments, and private interest groups to implement watershed plans that protect and restore native char.
- ! The Department will work within state laws and through local watershed planning processes to implement management actions needed to provide functioning habitats for native char defined in the WSP habitat Policy Framework including: 1) habitat protection and management; 2) basin hydrology and stream flow; 3) water sediment quality and sediment transport; 4) stream channel complexity; 5) riparian areas and wetlands; 6) lakes; and 7) marine areas.

## **Enforcement**

Enforcement objectives and implementation strategies are intended to support the population maintenance, fishery management, and habitat maintenance objectives. The primary purpose of the Enforcement Program is to protect the fish and wildlife resources of Washington State. The enforcement of fishing regulations, seasons and stream closures and hydraulic code regulations, will effectively meet enforcement objectives for protecting native char stocks.

**Enforce Seasons and Species-specific Fishing Regulations** - Regulations governing fishing in areas where native char occur will be enforced to ensure angler compliance.

*Implementation Strategy:*

- ! The Fish Management Program will coordinate with the Enforcement Program to identify and prioritize waters for enforcement emphasis activities. Data on violations will be reviewed periodically to assess angler compliance with regulations.

**Inform the Public Regarding Proper Identification of Native Char Species and Need for Their Protection.**

*Implementation Strategy:*

- ! Enforcement contacts with the public in the field will inform anglers of proper identification and conservation need for native char.
- ! Written and pictorial information will be distributed to the public to increase awareness of proper identification and conservation need for native char.

**Enforce Hydraulic Project Approvals** - Follow-up investigations of Hydraulic Project Approvals will be conducted to ensure permit compliance.

*Implementation Strategy:*

- ! Native char waters (including spawn timing and locations) will be distributed to each habitat and TFW biologist (Timber Fish and Wildlife biologist) responsible for processing HPAs or reviewing environmental documents ensure appropriate conditions are incorporated into these documents to protect native char habitat.
- ! Compliance checks will be made to insure adherence to conditions of permitted activities.

## **Research Monitoring and Evaluation**

The Department has identified below native char research questions which, when answered, will broaden the understanding of the ecological and taxonomic intricacies of these species, and will facilitate successful management efforts. Research objectives are identified that will fill critical information gaps important to the management of native char stocks. As new information is gathered and analyzed additional research questions will be prioritized for future investigation.

Native char stocks and habitats will be monitored to evaluate actions in meeting the objectives and goal for native char. This management plan will be dynamic, allowing for modifications of implementation strategies as identified through monitoring and evaluation. Investigative activities will be prioritized and implemented within Department budget.

**Determine Distribution of Native Char Stocks** - Standardized methodologies will be used to quantitatively determine the distribution of native char stocks in Washington.

*Implementation Strategy:*

- ! Standard methodologies identified in Bonar et al. (1997) will be used to collect and analyze distribution data. Distribution data will be maintained in a Geographic Information System database.

**Determine Stock Abundance and Biological Characteristics** - The status (abundance and natural biological characteristics) of native char stocks will be assessed and the bull trout and Dolly Varden salmonid stock inventory will be updated annually.

*Implementation Strategy:*

- ! Stock abundance information will be collected using methods presented in Bonar et al. (1997). Abundance data will be maintained in the Salmonid Stock Inventory database.
- ! Information on biological characteristics of native char stocks will be collected and reported in annual reports and maintained in the electronic databases.
- ! Abundance and biological data will be evaluated annually to update the status of bull trout and Dolly Varden stocks.

**Determine Genetic Diversity of Native Char Stocks** - Genetic analysis will be used to determine distribution of bull trout and Dolly Varden in Washington, identify their Major Ancestral Linages and Genetic Diversity Units, and identify the genetic relationships among the different native char life history types.

*Implementation Strategy:*

- ! Native char stocks will be sampled for genetic analysis using non-lethal means (fin clips).
- ! Fin tissue samples will be analyzed by the Department using microsatellite DNA techniques.

**Determine Length at Maturity and Frequency of Spawning for Individual Stocks** - The published literature indicates native char stocks may spawn in alternate years. Current fishery management strategies set minimum size limits that allow harvest on individuals only after the majority of females have had an opportunity to spawn at least once (minimum size limit are set at the upper end of the length frequency distribution for the age class containing the majority of first time spawning females).

*Implementation Strategy:*

- ! Determine reproductive frequency and length at maturity for stocks that currently have, or are expected to have, targeted harvest fisheries. Adjust minimum length criteria to ensure the age class containing the majority of females can spawn at least once prior to recruitment into the fishery.

**Determine Relationships Among Life History Types** - It is probable that there is some level of genetic exchange among the four bull trout and Dolly Varden life history forms. The extent of genetic exchange among resident, fluvial, adfluvial, and anadromous native char is largely unknown.

*Implementation Strategy:*

- ! Develop and implement research strategies to determine interactions between native char life history types.

**Monitor Fisheries** - The Department will monitor fisheries to ensure that direct and incidental harvest have no significant adverse impacts on long-term productivity of native char stocks.

*Implementation Strategy:*

- ! Native char harvest will be monitored by Department Fish and Wildlife Officers during routine and emphasis patrols.
- ! The Department may conduct creel surveys to estimate the harvest of native char in directed fisheries and incidentally caught in other target fisheries.
- ! The Department will determine the feasibility of expanding the current catch record card for recording harvest of bull trout and Dolly Varden.
- ! Conduct angler surveys periodically to determine anglers' knowledge, opinions, and preferences regarding native char management.

**Monitor Habitat Quality and Quantity** - Inventory and assess bull trout and Dolly Varden habitat periodically to evaluate changes in habitat quality and quantity over time.

*Implementation Strategy:*

- ! Conduct an inventory and assessment of native char habitat to evaluate changes in basin hydrology and stream flows, water and sediment quality and sediment transport, stream channel complexity, riparian and wetlands, lakes and reservoirs, marine areas and fish passage and access.

## Literature Cited

---

- Balon, E.K. 1984. Life Histories of Arctic Charrs: An Epigenetic Explanation of Their Invading Ability and Evolution. Pages 109-141 in L. Johnson and B.L. Burns, Editors. Biology of the Arctic Charr, Proceedings of the International Symposium on Arctic Charr, Winnipeg, Manitoba, May 1981. Univ. of Manitoba Press, Winnipeg.
- Baxter, J.S., E.B. Taylor, R.H. Devlin, J. Hagen, and J.K. McPhail. 1977. Evidence for natural hybridization between Dolly Varden (*Salvelinus malma*) and bull trout (*Salvelinus confluentus*) in a north central British Columbia watershed. Canadian Journal of Fisheries and Aquatic Science. 54:421-429.
- Behnke, R.J. 1980. A Systematic Review of the Genus *Salvelinus*. Pages 441-481 in E.K. Balon, Editor. CHARRS - Salmonid Fishes of the Genus *Salvelinus*. W. Junk Publishers, the Hague, the Netherlands.
- Bonar, S.A., M. Divens, and B. Bolding. 1997. Methods for sampling the distribution and abundance of bull trout/Dolly Varden. Washington Department of Fish and Wildlife Research Report No. RAD97-05. Olympia, WA. 48 p.
- Brown, L.G. 1992. On the Zoogeography and Life History of Washington's Native Char; Dolly Varden *Salvelinus malma* (Walbaum) and Bull Trout *Salvelinus confluentus* (Suckley). Report #94-04. Washington Department of Wildlife, Fisheries Management Division, Olympia, WA.
- Cavender, T.M. 1978. Taxonomy and Distribution of the Bull Trout, *Salvelinus confluentus* (Suckley), from the American Northwest. California Fish and Game, 3:139-174.
- Crane, P.A., L.W. Seeb, and J.E. Seeb. 1994. Genetic relationships among *Salvelinus* species inferred from allozyme data. Canadian Journal of Fisheries and Aquatic Science 51 (Supplement 1): 182-197.
- Johnston, T. and P. Mongillo. 1991. Unpublished char investigations data from files. Washington Department of Wildlife, Fisheries Management Division, Olympia, WA.
- Leary, R.F. and F.W. Allendorf. 1997. Genetic confirmation of sympatric bull trout and Dolly Varden in western Washington. Transactions of the American Fisheries Society 126:715-720.
- Kraemer, C. 1991. Unpublished char investigations data from files. Washington Department of Wildlife, Fisheries Management Division, Mill Creek, WA.
- Mongillo, P.E. 1993. The Distribution and Status of Bull Trout/Dolly Varden in Washington State. Report #93-22. Washington Department of Wildlife, Fisheries Management Division, Olympia, WA.

- Morton, W.M. 1970. On the Validity of All Subspecific Descriptions of North American *Salvelinus malma* (Walbaum). Copeia, 1970(3):581-587
- Savvaitova, K.A. 1980. Comments on the "Systematic Review of the Genus *Salvelinus*". Pages 480-481 in E.K. Balon, Editor. CHARRS - Salmonid Fishes of the Genus *Salvelinus*. W. Junk Publishers, the Hague, the Netherlands.
- Spruell, P. and F.W. Allendorf. 1997. Nuclear DNA analysis of Oregon bull trout Final Report to the Oregon Department of Fish and Wildlife Wild Trout and Salmon Genetics Laboratory Report 97/5. Division of Biological Sciences, University of Montana. Missoula Montana. 28 p.
- Utter, F. M. 1994. Current understanding of genetic population structure of bull trout. Unpublished report for Don Chapman Consultants, Boise, ID.
- Washington Department of Fish and Wildlife 1997a. Washington State salmonid stock inventory: bull trout/Dolly Varden. Washington Department of Fish and Wildlife. 428 p.
- Washington Department of Fish and Wildlife. 1997c. The watershed recovery inventory project report. Washington Department of Fish and Wildlife. Olympia, Washington. 10 p. + Appendices.
- Washington Department of Fisheries, Washington Department of Wildlife, and Western Washington Treaty Indian Tribes. 1993. 1992 Washington State Salmon and Steelhead Stock Inventory (SASSI). Mar. 1993. 212p.
- Washington Department of Game. 1984. A Basic Fishery Management Strategy for Resident and Anadromous Trout in the Stream Habitats of the State of Washington. Washington Department of Game, Fisheries Management Division, Olympia, WA.
- Washington Department of Wildlife. 1992. Draft Bull Trout/Dolly Varden Management and Recovery Plan. Washington Department of Wildlife, Fisheries Management Division, Olympia, WA. 125 p.
- Washington Fish and Wildlife Commission. 1995. Washington Department of Fish and Wildlife Goals, Policies, and Objectives by the Washington Fish and Wildlife Commission. Olympia, WA. 16 p.
- Washington Fish and Wildlife Commission. 1997a. Policy of Washington Department of Fish and Wildlife and Western Washington Treaty Tribes Concerning Wild Salmonids Adopted by Washington Fish and Wildlife Commission December 5, 1997. Olympia, WA. 46 p.
- Washington Fish and Wildlife Commission. 1997b. Additional Policy Guidance on Deferred Issues Concerning Wild Salmonid Policy Adopted by Washington Fish and Wildlife Commission December 5, 1997. Olympia, WA. 21 p.

Wright, S. 1992. Guidelines for selecting regulations to manage open-access fisheries for natural populations of anadromous and resident trout in stream habitats. North American Journal of Fisheries Management 12: 517-527



# Glossary

---

**Critical Stock:** A stock of fish experiencing production levels that are so low that permanent damage to the stock is likely or has already occurred.

**Depressed Stock:** A stock of fish whose production is below expected levels based on available habitat and natural variations in survival levels, but above the level where permanent damage to the stock is likely.

**Genetic Diversity Unit:** A group of genetically similar stocks that is genetically distinct from other such groups. The stocks typically exhibit similar life histories and occupy ecologically, geographically, and geologically similar habitats. A GDU may consist of a single stock.

**Hybridization:** The genetic interbreeding of fish from two or more different stocks or species (e.g., brook trout interbreeding with bull trout).

**Native Species:** A species of fish indigenous to Washington State.

**Stock:** The fish spawning in a particular lake or stream(s) (or portion of it) at a particular season, which to a substantial degree, do not interbreed with any group spawning in a different place, or in the same place at a different season.

**Healthy Stock:** A stock of fish experiencing production levels consistent with its available habitat and within the natural variations in survival for the stock.

**Unknown Stock:** This description is applied to stocks where there is insufficient information to identify stock status with confidence.